



© 1995-2012 by Isidoro Martínez

Properties of gases

(Function values at 100 kPa and 288 K or the normal boiling temperature if greater)

| Substance | Formula | Molar mass | Boiling temp. | Critical temp | Critical pressure | Critical compressibility factor ^a | Pitzer's acentric factor | Thermal capacity ^b | Thermal conductivity ^c | Dynamic viscosity ^d |
|-------------------------------|---|-------------|--------------------|------------------|-------------------|--|--------------------------|-------------------------------|-----------------------------------|--------------------------------|
| | | M kg/mol | T_b K | T_{cr} K | p_{cr} MPa | Z_{cr} | ω | c_p J/(kg·K) | k W/(m·K) | $\mu \cdot 10^6$ Pa·s |
| Acetone | C ₃ H ₆ O | 0.058 | 329.2 | 508 | 4.70 | 0.233 | 0.309 | 1300 | | |
| Acetylene | C ₂ H ₂ | 0.026 | 189.5 ^f | 309 | 6.20 | 0.271 | 0.184 | 1580 | 0.019 | 9.3 |
| Air | 79%N ₂ ⁺ 21%O ₂ | 0.029 | 82 ^e | 132 ^g | 3.75 ^g | 0.28 ^g | 0.035 | 1004 | 0.024 | 18.1 |
| Ammonia | NH ₃ | 0.017 | 239.8 | 406 | 11.30 | 0.242 | 0.250 | 2200 | 0.022 | 9.3 |
| Argon | Ar | 0.040 | 87.4 | 151 | 4.86 | 0.291 | 0 | 523 | 0.016 | 21.0 |
| Benzene | C ₆ H ₆ | 0.078 | 353.3 | 563 | 4.92 | 0.271 | 0.212 | 1300 | 0.007 | 7.0 |
| 1,3-Butadiene | C ₄ H ₆ | 0.054 | 268.5 | 425 | 4.33 | 0.270 | 0.193 | 1510 | | |
| n-Butane | C ₄ H ₁₀ | 0.058 | 272.6 | 425 | 3.80 | 0.274 | 0.193 | 1580 | 0.015 | 7.0 |
| iso-Butane | C ₄ H ₁₀ | 0.058 | 261.5 | 408 | 3.64 | 0.280 | 0.176 | 1580 | 0.015 | 9.0 |
| Carbon dioxide | CO ₂ | 0.044 | 194.7 ^f | 304 | 7.38 | 0.274 | 0.225 | 840 ^h | 0.015 | 9.0 |
| Carbon monoxide | CO | 0.028 | 81.7 | 133 | 3.50 | 0.295 | 0.049 | 1100 | 0.023 | 17.0 |
| Carbon tetrachloride | CCl ₄ | 0.154 | 349.7 | 556 | 4.56 | 0.272 | 0.194 | 862 | 0.017 | 16.0 |
| Cyclohexane | C ₆ H ₆ | 0.084 | 353.9 | 554 | 4.07 | 0.273 | 0.212 | | | |
| n-Decane | C ₁₀ H ₂₂ | 0.142 | 447.3 | 619 | 2.12 | 0.247 | 0.490 | 1680 | | |
| n-Dodecane | C ₁₂ H ₂₆ | 0.170 | 489.4 | 659 | 1.80 | 0.240 | 0.562 | 1690 | | |
| DME (dimethyl ether) | C ₂ H ₆ O | 0.046 | 250.6 | 400 | 5.37 | 0.271 | 0.274 | 1430 | | |
| Ethane | C ₂ H ₆ | 0.030 | 184.6 | 305 | 4.88 | 0.285 | 0.100 | 1700 | 0.020 | 11.0 |
| Ethanol | C ₂ H ₆ O | 0.046 | 351.7 | 516 | 6.39 | 0.248 | 0.635 | 1520 | 0.013 | 14.2 |
| Ether (diethyl ether) | C ₄ H ₁₀ O | 0.074 | 307.6 | 467 | 3.61 | 0.260 | 0.281 | 1600 | 0.015 | 7.5 |
| ETBE (ethyl tert-butyl ether) | C ₆ H ₁₄ O | 0.102 | 345 | 517 | 3.11 | 0.274 | 0.298 | 1550 | | |
| Ethylene | C ₂ H ₄ | 0.028 | 169.5 | 283 | 5.12 | 0.276 | 0.085 | 1470 | 0.018 | 9.6 |
| Ethylene glycol | C ₂ H ₆ O ₂ | 0.062 | 471 | 645 | 7.53 | 0.268 | 1.137 | 1410 | | |
| Helium (⁴ He) | He | 0.004 | 4.2 | 5.3 | 0.23 | 0.301 | -0.387 | 5190 | 0.142 | 19.0 |
| Helium 3 (³ He) | He | 0.003 | 1 | 3.3 | 0.11 | 0.301 | -0.460 | | | |
| n-Heptane | C ₇ H ₁₆ | 0.100 | 371 | 540 | 2.77 | 0.263 | 0.350 | 1650 | 0.013 | 6.5 |
| n-Hexane | C ₆ H ₁₄ | 0.086 | 342 | 508 | 3.03 | 0.263 | 0.296 | 1700 | 0.014 | 6.5 |

| | | | | | | | | | | |
|--|--|-------|--------------------|-------|-------|-------|-------|-------|-------|------|
| Hydrazine | N ₂ H ₄ | 0.032 | 387 | 653 | 14.7 | | 0.315 | | | |
| Hydrogen | H ₂ | 0.002 | 20.1 | 33 | 1.32 | 0.305 | -0.22 | 14200 | 0.168 | 8.4 |
| (Hydrogen) Deuterium | D ₂ | 0.004 | 23.6 | 38 | 1.66 | 0.249 | -0.16 | 14200 | 0.131 | 12 |
| Mercury ⁱ | Hg | 0.201 | 630 | 736 | 104 | | | | | |
| Methane | CH ₄ | 0.016 | 112 | 191 | 4.60 | 0.288 | 0.010 | 2180 | 0.031 | 10.3 |
| Methanol | CH ₄ O | 0.032 | 338.1 | 513 | 8.08 | 0.224 | 0.559 | 1350 | 0.015 | 9.8 |
| MTBE (methyl tert-butyl ether) | C ₅ H ₁₂ O | 0.088 | 328 | 497 | 3.43 | 0.273 | 0.267 | 1500 | | |
| Neon | Ne | 0.020 | 26.2 | 44 | 2.70 | 0.301 | 0 | 1030 | 0.046 | 30.0 |
| Nitrogen | N ₂ | 0.028 | 77.4 | 126 | 3.39 | 0.290 | 0.038 | 1040 | 0.024 | 16.6 |
| Nitrogen dioxide ^j | NO ₂ | 0.046 | 294.5 | 431 | 10.1 | | | 783 | 0.017 | 130 |
| Nitrogen monoxide | NO | 0.030 | 121.2 | 180 | 6.55 | 0.250 | 0.607 | 996 | 0.024 | 29.4 |
| di-Nitrogen oxide ^k | N ₂ O | 0.044 | 184.7 | 310 | 7.26 | 0.272 | 0.141 | 864 | 0.015 | 13.6 |
| n-Octane | C ₈ H ₁₈ | 0.114 | 399.1 | 569 | 2.49 | 0.259 | 0.394 | 1700 | 0.011 | 5.5 |
| Ozone | O ₃ | 0.048 | 161.4 | 268 | 6.78 | 0.272 | | | | |
| Oxygen | O ₂ | 0.032 | 90.2 | 155 | 5.08 | 0.288 | 0.021 | 913 | 0.024 | 19.1 |
| iso-Pentane | C ₅ H ₁₂ | 0.072 | 301.3 | 461 | 3.33 | 0.268 | 0.227 | 1680 | 0.015 | 11.7 |
| n-Pentane | C ₅ H ₁₂ | 0.072 | 309.2 | 470 | 3.38 | 0.262 | 0.251 | 1680 | 0.015 | 11.7 |
| Phenol | C ₆ H ₆ O | 0.094 | 455 | 694 | 6.13 | 0.243 | 0.426 | | | |
| Propane | C ₃ H ₈ | 0.044 | 231.1 | 370 | 4.26 | 0.281 | 0.152 | 1570 | 0.015 | 7.4 |
| iso-Propanol | C ₃ H ₈ O | 0.060 | 355.4 | 508 | 4.76 | 0.248 | 0.669 | | | |
| Propylene (propene) | C ₃ H ₆ | 0.042 | 225.4 | 365 | 4.62 | 0.275 | 0.148 | 1460 | 0.014 | 8.1 |
| Propylene glycol | C ₃ H ₈ O ₂ | 0.076 | 461.3 | 626 | 6.10 | 0.280 | 1.107 | | | |
| R-12 (CFC-12) (dichlorodifluoromethane) | CCl ₂ F ₂ | 0.121 | 243.0 | 385 | 4.14 | 0.280 | 0.176 | 573 | 0.008 | 12.5 |
| R-134a (HFC-134a) (tetrafluoroethane) | CF ₃ CH ₂ F | 0.102 | 246.6 | 374 | 4.07 | 0.258 | 0.332 | 840 | 0.014 | 12.2 |
| Sulfur dioxide | SO ₂ | 0.064 | 263.2 | 430 | 7.87 | 0.264 | 0.251 | 607 | 0.009 | 11.6 |
| Sulfur hexafluoride ^l | SF ₆ | 0.146 | 204.9 ^f | 319 | 3.76 | 0.360 | 0.210 | 598 | 0.12 | 16.0 |
| Toluene | C ₇ H ₈ | 0.092 | 383.7 | 592 | 4.13 | 0.284 | 0.266 | | | |
| Uranium hexafluoride ^m | UF ₆ | 0.352 | 329 ^f | 503 | 4.60 | 0.282 | 0.092 | 370 | 0.009 | 20 |
| Water ⁿ | H ₂ O | 0.018 | 372.8 | 647.3 | 22.12 | 0.229 | 0.344 | 2050 | 0.025 | 12.1 |

^aCritical molar volumes can be obtained from $v_{cr} = Z_{cr}RT_{cr}/p_{cr}$, and critical densities from $\rho_{cr} = M/v_{cr}$ (e.g. for acetone $v_{cr} = 209 \cdot 10^{-6} \text{ m}^3/\text{mol}$ and $\rho_{cr} = 351 \text{ kg/m}^3$).

^bThermal capacities of monoatomic gases do not change with temperature, but for polyatomic gases it increases more the

more atoms has the molecule.

^c Thermal conductivity of gases increases with the square root of temperature, and do not change with pressure. Thermal diffusivity $a \equiv k / (\rho c_p)$. According to simple generalised transport theory in gases, thermal diffusivity, mass diffusivity and kinematic viscosity of gases have the same values.

^d Dynamic viscosity of gases increases with the square root of temperature, and do not change with pressure. Kinematic viscosity $\nu \equiv \mu / \rho$.

^e Bubble point.

^f Sublimation point.

^g Pseudo-critical point (Kay's model).

^h Most gas properties vary a lot near the critical point, what may be here the case; e.g., for CO₂ gas at 288 K and 100 kPa, thermal capacity at constant pressure is $c_p = 840$ J/(kg·K), growing at constant $T = 288$ K from $c_p = 833$ J/(kg·K) at very low pressure, to $c_p = 3010$ J/(kg·K) at the saturation pressure (5063 kPa). Thermal capacity in the ideal gas limit ($p \rightarrow 0$) varies almost linearly (e.g. $c_p = 753$ J/(kg·K) at the triple-point temperature, $c_p = 850$ J/(kg·K) at the critical-point temperature).

ⁱ Mercury is obtained by oxidation of cinnabar at some 600 °C and vapour condensation. Mercury vapour should not exceed 0.1 mg/m³ in breathing air (notice that saturated air at 20 °C already contains more than that limit).

^j Nitrogen dioxide, NO₂, is a very toxic brown gas at normal conditions (but readily condensable, $T_b = 11$ °C). All nitrogen oxides slowly decomposing to nitrogen and oxygen, making it difficult to keep them in pure state; besides, NO₂ readily dimerises to dinitrogen tetroxide, N₂O₄, a colourless gas with double density than NO₂ (e.g. when heating from above an ampoule containing NO₂, some N₂O₄ is formed at the top ($2\text{NO}_2(\text{g}) = \text{N}_2\text{O}_4(\text{g}) + 57$ kJ/mol), which can be seen sinking to the bottom because of buoyancy).

^k Di-nitrogen oxide, N₂O, also known as nitrous oxide (NO is nitric oxide), or nitrogen hemi-oxide, or nitrogen protoxide, or laughing gas, is used in respiratory anaesthesia since the pioneering trials of Sir Humphrey Davy in 1789 shortly after its discovery by J. Priestley in 1772, as a non-flammable non-ozone-depleting propellant in aerosol cans, and as a fuel additive to enhance combustion (it liberates oxygen; if added as compressed liquid in the intake manifold, it greatly increases fuel load). It has a global warming potential (GWP) of 300 times that of CO₂, being the third contributor to anthropogenic GWP, after CO₂ and CH₄.

^l Sulfur hexafluoride is a synthetic gas used as insulator for electrical equipment (breakdown potential three times larger than air, and as a fluorine source for edging in the electronics industry. It is a non-flammable, non toxic gas, which decomposes at 750 K; it has low water solubility, and a very large IR absorbance (it is the most potent greenhouse gas, GWP=22 000), what has been used as a trace gas for gas-leakage detection.

^m Uranium hexafluoride, perhaps the heaviest simple molecule, is the only uranium compound presently used in industrially enrichment of U-235, both on gas diffusion and on gas centrifugation processes. At room conditions, it is a white crystalline solid with a high vapour pressure ($p_v = 11$ kPa at 20 °C).

ⁿ The boiling point of water at 100 kPa is $T_b = 372.75 \pm 0.02$ K (99.60 ± 0.02 °C), whereas at 101.325 kPa (1 atm) it is 373.12 ± 0.02 K (99.97 ± 0.02 °C).